



## Associations of plasma CD36 and body fat distribution

Wang, Yeli; Koch, Manja; di Giuseppe, Romina; Evans, Kirsten; Borggrefe, Jan; Nöthlings, Ute; Handberg, Aase; Jensen, Majken K; Lieb, Wolfgang

*Published in:*  
Journal of Clinical Endocrinology and Metabolism

*DOI (link to publication from Publisher):*  
[10.1210/jc.2019-00368](https://doi.org/10.1210/jc.2019-00368)

*Publication date:*  
2019

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*  
Wang, Y., Koch, M., di Giuseppe, R., Evans, K., Borggrefe, J., Nöthlings, U., Handberg, A., Jensen, M. K., & Lieb, W. (2019). Associations of plasma CD36 and body fat distribution. *Journal of Clinical Endocrinology and Metabolism*, 104(9), 4016-4023. <https://doi.org/10.1210/jc.2019-00368>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

## Associations of plasma CD36 and body fat distribution

Yeli Wang, Manja Koch, Romina di Giuseppe, Kirsten Evans, Jan Borggrefe, Ute Nöthlings, Aase Handberg, Majken K. Jensen, Wolfgang Lieb

*The Journal of Clinical Endocrinology & Metabolism*  
Endocrine Society

Submitted: February 13, 2019

Accepted: April 23, 2019

First Online: April 29, 2019

Advance Articles are PDF versions of manuscripts that have been peer reviewed and accepted but not yet copyedited. The manuscripts are published online as soon as possible after acceptance and before the copyedited, typeset articles are published. They are posted "as is" (i.e., as submitted by the authors at the modification stage), and do not reflect editorial changes. No corrections/changes to the PDF manuscripts are accepted. Accordingly, there likely will be differences between the Advance Article manuscripts and the final, typeset articles. The manuscripts remain listed on the Advance Article page until the final, typeset articles are posted. At that point, the manuscripts are removed from the Advance Article page.

**DISCLAIMER:** These manuscripts are provided "as is" without warranty of any kind, either express or particular purpose, or non-infringement. Changes will be made to these manuscripts before publication. Review and/or use or reliance on these materials is at the discretion and risk of the reader/user. In no event shall the Endocrine Society be liable for damages of any kind arising references to, products or publications do not imply endorsement of that product or publication.

## Plasma CD36 and body fat distribution

**Associations of plasma CD36 and body fat distribution**

Yeli Wang<sup>1\*</sup>, Manja Koch<sup>2\*</sup>, Romina di Giuseppe<sup>3</sup>, Kirsten Evans<sup>2</sup>, Jan Borggrefe<sup>4</sup>, Ute Nöthlings<sup>5</sup>, Aase Handberg<sup>6,7</sup>, Majken K. Jensen<sup>2</sup>, Wolfgang Lieb<sup>3</sup>

<sup>1</sup>Health Services and Systems Research, Duke-NUS Medical School, Singapore;

<sup>2</sup>Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA, USA;

<sup>3</sup>Institute of Epidemiology, Kiel University, Kiel, Germany;

<sup>4</sup>Department of Neuroradiology, University Hospital Cologne, Cologne, Germany;

<sup>5</sup>Department of Nutrition and Food Sciences, University of Bonn, Bonn, Germany;

<sup>6</sup>Department of Clinical Biochemistry, Aalborg University Hospital, Aalborg, Denmark;

<sup>7</sup>Department of Clinical Medicine, The Faculty of Medicine, Aalborg University, Aalborg, Denmark;

**ORCID numbers:**

0000-0003-3031-6199

Wang

Yeli

Received 13 February 2019. Accepted 23 April 2019.

\*The authors contributed equally to the paper.

**Context:** CD36 is a class B scavenger-receptor involved in the uptake of fatty acids in liver and adipose tissue. It is unknown whether plasma CD36 levels are related to liver fat content or adipose tissue in the general population.

**Methods:** We measured plasma CD36 from 575 participants of the community-based PopGen-cohort who underwent magnetic resonance imaging (MRI) to quantify visceral (VAT) and subcutaneous (SAT) adipose tissue and liver signal intensity (LSI), a proxy for liver fat content. Non-alcoholic fatty liver disease (NAFLD) was defined as LSI  $\geq 3.0$  in the absence of high alcohol intake. The relations between plasma CD36 and body mass index (BMI), VAT, SAT, LSI, and NAFLD were evaluated using multivariable-adjusted linear and logistic regression analysis.

**Results:** Plasma CD36 concentrations were correlated with BMI ( $r=0.11$ ;  $P=0.01$ ), SAT ( $r=0.16$ ;  $P<0.001$ ), and VAT ( $r=0.15$ ,  $P<0.001$ ), but not with LSI ( $P=0.44$ ). In multivariable-adjusted regression models, mean BMI values rose across CD36-quartiles (Q1: 27.8 kg/m<sup>2</sup>; Q4: 28.9 kg/m<sup>2</sup>;  $P$ -trend=0.013). Similarly, VAT (Q1: 4.13 dm<sup>3</sup>; Q4: 4.71 dm<sup>3</sup>;  $P$ -trend<0.001) and SAT (Q1: 7.61 dm<sup>3</sup>; Q4: 8.74 dm<sup>3</sup>;  $P$ -trend<0.001) rose across CD36 quartiles. Plasma CD36 concentrations were unrelated to LSI ( $P$ -trend=0.36), and NAFLD ( $P$ -trend=0.64). Participants with NAFLD and elevated alanine aminotransferase (ALT), a marker for liver damage, had higher CD36 compared to NAFLD participants with normal ALT.

**Conclusions:** Higher plasma concentrations of CD36 were associated with greater general and abdominal adiposity, but not with liver fat content or NAFLD in this community-based sample. However, plasma CD36 may reflect more severe liver damage in NAFLD.

We provide evidence that plasma CD36 levels are associated with general and abdominal adiposity, but not with MRI-determined liver fat content, in a general population sample. .

## INTRODUCTION

The global prevalence of obesity has tripled in the past two decades (1). In 2016, more than 650 million adults worldwide were obese, representing 13% of the total global population (1). Up to 90% of obese patients have non-alcoholic fatty liver disease (NAFLD) (2), which represents the most common chronic liver condition, characterized by liver fat accumulation (3). Both obesity and NAFLD are significant risk factors for increased morbidity and mortality, e.g. from type 2 diabetes, cardiovascular disease and cancer (4,5). Therefore, it is of scientific importance to deepen our understanding of the underlying etiology and look for effective targets for the prevention and treatment of obesity.

CD36 is a multifunctional glycosylated protein widely expressed in a variety of tissues (6). In metabolically active tissues such as liver and adipose tissue, CD36 functions as a lipid transporter that binds and uptakes long-chain fatty acids and thereby fosters fat accumulation in hepatocytes and adipocytes (6). In line with its biological function, CD36 expression has been reported to be upregulated in hepatocytes of patients with fatty liver (7,8), and in adipose tissue of obese individuals (9). In addition, human genetic studies have identified several genetic variants in the *CD36* gene to be associated with traits reflecting lipid metabolism and obesity (10-12).

Traditionally, the measurement of CD36 required fresh blood or tissue samples and was therefore not feasible in large-scale population studies. Recent research efforts enabled the assessment of plasma CD36 in stored plasma samples (13). In subsequent studies, plasma CD36 levels were correlated with CD36 expression (14) and with body mass index (BMI) and other indices of body fat distribution (13,15-19). In addition, positive correlations have been observed between plasma CD36 levels and liver fat in patients with higher liver fat content (NAFLD and chronic hepatitis C) (15,20,21) and obesity (17,18). However, very little is known about the association of plasma CD36 levels with liver fat and other fat depots in the general population. In 1296 healthy individuals free of diabetes and hypertension, a positive association of plasma CD36 levels with two indices of fatty liver has been reported (22). In that study, however, no direct measure of liver fat was available; instead, liver fat was estimated using two algorithms based on anthropometric (BMI, waist circumference [WC]) and biochemical (alanine transaminase [ALT], gamma-glutamyl transferase and aspartate aminotransferase, levels of triglycerides and insulin) measures, as well as on the presence of metabolic syndrome and type 2 diabetes (22).

Non-invasive imaging techniques quantify abdominal fat depots and liver fat content more accurately than surrogate anthropometrical measures including WC, BMI or liver fat algorithms (23). Therefore, in the present analysis, we aimed to relate plasma CD36 concentrations in cross-sectional analyses to magnetic resonance imaging (MRI)-traits of liver fat content and body fat distribution in a community-based sample from Northern Germany (24). Since the expression of tissue CD36 is much higher in adipose tissue than liver (25), we hypothesize that plasma CD36 levels are more strongly related to visceral adipose tissue than liver fat content in a general population.

## METHODS

### Study sample

The PopGen control cohort is an ongoing prospective cohort study among a random sample of 1316 individuals living in northern Germany. The design has been described in detail previously (26). In brief, at baseline recruitment between 2005 and 2007, the PopGen biobank randomly identified 23,000 local residents through population registries in the city of Kiel and invited them to participate in the study. A total of 4,267 participants agreed to take part, and among them, 747 subjects agreed to participate in the follow-up study. In addition,

the PopGen biobank recruited 569 blood donors at the University Hospital Schleswig-Holstein in Kiel, and these two groups constituted the final PopGen control cohort (1316 individuals) (27). The first follow-up examination was conducted between 2010 and 2012, and 952 participants agreed to be re-examined at the study center. During this follow-up examination, trained study nurses performed a medical examination and drew blood. Furthermore, a whole-body MRI was part of this examination cycle, including an assessment of liver fat content and body fat distribution. Information on demographics, dietary intake, lifestyle factors and medical history was collected via self-administered questionnaires (24,26,28).

For the present analysis, we used data from the first follow-up examination. A total of 575 participants underwent MRI to assess VAT and SAT, as well as liver fat content. The flowchart of the current study design is shown in **Supplemental Figure S1**. The study procedures were approved by the ethical review board of the Medical Faculty of Kiel University, Germany. All study participants provided written informed consents.

### Assessment of covariates

Information on participants' characteristics such as age, sex, education, smoking status, alcohol consumption, physical activity and medical history was collected using self-administered questionnaires. Alcohol consumption during the past year was calculated as the means of the reported values from a validated food frequency questionnaire for German populations (29). We quantified physical activity as weekly metabolic equivalent task (MET)-hours by summing the reported weekly hours in different activities (walking, cycling, sports, gardening, do-it-yourself activities and household tasks (28)) and multiplying them by their corresponding MET values averaged for winter and summer (30). Prevalent type 2 diabetes was determined by self-reported physician-diagnosed type 2 diabetes, antidiabetic medication usage, fasting blood glucose levels  $\geq 126$  mg/dL or hemoglobin A1c levels  $\geq 6.5\%$  at the baseline or at the follow-up visit. In addition, prevalent hypertension was defined as self-reported hypertension diagnosed by a physician, using hypertension medication, or having diastolic blood pressure  $\geq 90$  mm Hg or systolic blood pressure  $\geq 140$  mm Hg (24,28).

Trained study nurses measured WC, body weight and height of participants wearing light clothing without shoes (28,31). Study nurses subtracted 2 kilograms from the measured weight to correct for clothing. BMI was calculated as weight (in kilograms) divided by height (in meters) squared. WC was measured at the midpoint between the spina iliaca crest and lower costal margin (31). Study nurses measured blood pressure three times by sphygmomanometry after the participant rested for at least 5 minutes. We calculated the arithmetic means of the last two blood pressure measurements (28).

### Biochemical measurements

All participants provided blood samples in a sitting position after overnight fasting. Most biosamples were stored at -80 degrees until biomarkers were assayed.

Standard laboratory parameters (including concentrations of total cholesterol [TC], triglycerides [TG], high-density lipoprotein cholesterol [HDL-C], low-density lipoprotein cholesterol [LDL-C]) were measured on the day of the examination at the Institute for Clinical Chemistry, University Hospital Schleswig-Holstein, Kiel, Germany (28). ALT was assessed by enzymatic colorimetry (Hitachi Modular; Roche Diagnostics), C-reactive protein (CRP) levels were determined by immunoturbidimetry, blood glucose levels by enzymatic ultraviolet tests (Roche Diagnostics) and HbA1c levels by HPLC and photometric detection (Bio-Rad Laboratories). Plasma CD36 levels were assayed (from samples stored at -80 degrees) at Aarhus University Hospital by an *in-house* ELISA assay (13), and the intra-assay coefficient of variation was 11%.

### Outcome assessment



We used MRI to quantify the liver fat content and adipose tissue volumes on a Magnetom Avanto 1.5-T whole-body imager (Siemens Medical Solutions) (28,32). Study nurses performed MRI with participants at the supine position with arms extended over the head (32). In-phase and out-of-phase images were acquired during a breath hold by using axial T1-weighted gradient echo MRI with a 10.4-ms repetition time, echo time of 4.76 ms (in-phase) and 7.14 ms (opposed phase), 10° flip angle, 128 × 80 matrix, and 275 × 440-mm field of view (24). Liver fat content was quantified as the relative liver signal intensity (LSI) difference of the liver on out-of-phase compared with in-phase images in arbitrary units (24,33). We validated the accuracy of LSI to quantify liver fat content using T2-corrected <sup>1</sup>H single-voxel proton magnetic resonance spectroscopy (MRS) on a 1.5-T whole-body imager (Siemens Medical Solutions) as the reference method in a sample of 40 individuals (55% men; median age: 55.6 y; median BMI: 27.7 kg/m<sup>2</sup>) (24). Spearman's correlation coefficient between LSI values and the MRS-determined liver fat content was  $r = 0.609$  ( $P < 0.001$ ) (24). Fatty liver disease (FLD) was determined as log-transformed LSI  $\geq 3.0$  according to a cut-off value determined by receiver operating characteristic (ROC) analysis using spectroscopic-determined FLD (liver fat  $\geq 5.56\%$ ) as the reference (area under ROC curve: 0.86; 95% confidence interval [CI], 0.74-0.98) (24,34). The log-LSI cut-off of  $\geq 3.0$  was identified by Youden Index, which corresponded to a sensitivity of 0.79 and a specificity of 0.81 (24). NAFLD was defined as having FLD in the absence of higher alcohol intake defined as  $\leq 7$  drinks per week for women and  $\leq 14$  drinks per week for men (35,36).

In addition, VAT was determined as the sum of VAT voxels from the top of the liver to the femoral heads inside the abdominal muscular wall, and SAT was determined as the sum of adipose tissue voxels underneath the skin layer surrounding the abdomen from the top of the liver to the femur heads (32). Both VAT and SAT were expressed in decimeters cubed (dm<sup>3</sup>) (28).

### Statistical analysis

To describe the characteristics of the study participants, we divided the study sample into quartiles based on the distribution of plasma CD36 concentrations using sex-specific cut-offs. Continuous variables with normal distribution were reported as mean  $\pm$  standard deviation, while continuous variables with skewed distribution were reported as median (Q1, Q3), and categorical variables were presented as percentages.

Spearman partial correlations adjusted for age and sex were estimated between plasma CD36 concentrations and LSI, adiposity measures (BMI, WC, VAT, SAT) and blood biomarkers, including ALT, glucose, HbA1c, TC, TG, HDL-C, LDL-C, CRP and fetuin-A. For participants ( $n=229$ ) whose CRP values were below the detection limit (0.9 mg/L), half of the detection limit (0.45 mg/L) was assigned as their CRP values.

Cubic spline analysis with four knots at the 5<sup>th</sup>, 35<sup>th</sup>, 65<sup>th</sup> and 95<sup>th</sup> percentiles (50<sup>th</sup> percentile as referent) was used to assess potential non-linearity of the relations between plasma CD36 and FLD as well as NAFLD. When assessing the association between plasma CD36 and NAFLD, we excluded participants with self-reported liver disease (hepatitis A, B, C, or D virus infection, hemochromatosis, autoimmune liver disease, or liver cirrhosis,  $n=18$ ) other than FLD. To analyze the associations between plasma CD36 and FLD as well as NAFLD, we used multivariable-adjusted logistic regression models and calculated odds ratios for FLD using the lowest plasma CD36 quartile as the reference category. We adjusted for sex, age (continuous), years of education [ $\leq 9$  years (no level of general education completed or secondary general school-leaving certificate), 10 years (intermediate school-leaving certificate), or  $\geq 11$  years (university of applied sciences or university entrance qualification)], smoking status [never (smoking period  $\leq 3$  months), former (smoking period  $> 3$  months), or current], physical activity (metabolic equivalent task hours/week; continuous), alcohol consumption (continuous) and history of diabetes (yes or no) as potential

confounding factors. We additionally adjusted for BMI (model 2) or VAT/SAT (model 3) to assess whether the association of CD36 with FLD and NAFLD is independent of these adiposity measures. Adjusted means of LSI, BMI, VAT and SAT according to CD36 quartiles were calculated by analysis of covariance.

Potential effect modification by sex, median age (< vs.  $\geq 62.3$  years), prevalent diabetes and prevalent hypertension were tested by including respective multiplicative interaction terms in the fully-adjusted regression models.

Analyses were performed with SAS software, version 9.4 (SAS Institute, Cary, North Carolina). All tests were two-sided, and *P* values <0.05 were considered to be statistically significant.

## RESULTS

The median age of this sample was 62.3 (range: 25.1–82.8) years at the first follow-up examination. Characteristics of the study sample according to the sex-specific quartiles of plasma CD36 levels are shown in **Table 1**. In unadjusted statistical models, compared to people with lower plasma CD36 levels, individuals with higher CD36 levels had higher mean values for different adiposity measures (BMI, WC, SAT, VAT), diastolic and systolic blood pressure, ALT, a higher prevalence of FLD and NAFLD and lower education levels.

Plasma CD36 levels were positively correlated with BMI, WC, SAT, VAT, ALT, total cholesterol and TG but not with LSI ( $r = 0.03$ ;  $P = 0.44$ ), HDL-C ( $r = -0.01$ ;  $P = 0.85$ ), or LDL-C ( $r = 0.07$ ;  $P = 0.09$ ) in age- and sex-adjusted analyses (**Supplemental Table S1**).

In multivariable-adjusted least square-means analyses using the first CD36-quartile as referent, participants in the second, third and fourth CD36 quartile had significantly higher levels of BMI, SAT and VAT with a graded increase across quartiles (all *P*-trend <0.05; **Table 2**), however, such a graded association was not observed for LSI (*P*-trend >0.10). In addition, BMI, SAT and VAT explained 8%, 17% and 13% of the variance in plasma CD36 concentrations, respectively; whereas LSI explained 1% of the variance of plasma CD36.

The association of plasma CD36 levels with FLD and NAFLD is shown in **Supplemental Table S2**. Restricted cubic spline analysis did not suggest a nonlinear association of plasma CD36 levels with FLD and NAFLD (data not shown). After adjusting for age, sex, education levels, smoking status, alcohol intake, physical activity and prevalent diabetes, higher CD36 concentrations conferred slightly higher odds for FLD (highest vs. lowest quartile, odds ratio [OR]: 1.13; 95% CI: 0.69, 1.87) and NAFLD (highest vs. lowest quartile, OR: 1.28; 95% CI: 0.74, 2.22), but the respective odds ratios were not statistically significant. The risk estimate of CD36 for liver fat was further attenuated towards the null after adjustment for BMI (highest vs. lowest quartile, OR for FLD: 0.98; 95% CI: 0.58, 1.64; OR for NAFLD: 1.09; 95% CI: 0.62, 1.92) or adjustment for VAT/SAT (highest vs. lowest quartile, OR for FLD: 1.02; 95% CI: 0.61, 1.70; OR for NAFLD: 1.15; 95% CI: 0.66, 2.01; **Supplemental Table S2**).

We also compared mean CD36 levels in individuals with NAFLD and normal ALT concentrations ( $\leq 33$  IU/L in men and  $\leq 25$  IU/L in women;  $n = 113$ ) to CD36 levels of participants with NAFLD and elevated ALT concentrations ( $n = 70$ ). CD36 levels were statistically significantly higher in individuals with NAFLD and ALT ( $1.58 \pm 0.82$  units vs.  $1.21 \pm 0.72$  units;  $P = 0.002$ ).

We found no evidence for effect modification by sex, age, prevalent diabetes and prevalent hypertension (all *P*-interaction >0.05). All supplemental materials were stored at the public data repository (37).

## DISCUSSION

### Principal observations

In this cross-sectional analysis in a population-based sample, we observed statistically significant correlations between plasma CD36 levels and different markers of general and abdominal adiposity, including BMI, WC, VAT and SAT. By contrast, no association of plasma CD36 concentrations with liver fat content, FLD and NAFLD was observed.

#### **In the context of the published literature**

With the present analyses on MRI-derived traits of liver fat, SAT and VAT in addition to BMI and WC, we expand on prior studies that have reported positive correlations of plasma CD36 with adiposity markers (mainly BMI and WC) (13,15,17-19,22) and liver fat content (15,17,18,20-22) in other European samples, mainly in samples with defined disease conditions such as liver disease or obesity. Several prior studies have consistently observed a positive correlation between plasma CD36 levels and BMI (13,17,18,22). However, since BMI does not differentiate between body fat and muscle weight and since visceral fat may play a greater role in contributing to metabolic diseases than subcutaneous fat (38,39), more detailed analyses regarding the association of CD36 levels with different fat depots were warranted. In 111 NAFLD patients and 33 healthy controls, an inverse correlation between plasma CD36 levels and dual-energy X-ray absorbiometry VAT ( $r=-0.21$ ;  $P<0.05$ ), but no correlation with SAT, were reported (15). Other indicators of body composition such as WC, fat mass, central fat mass and truncal fat mass have also shown to be positively correlated with plasma CD36 levels in women with polycystic ovary syndrome (19), in obese children (18) and in morbidly obese individuals (17) receiving lifestyle intervention programs and gastric bypass surgeries, respectively, for weight loss.

In our comprehensive analyses in a community-based sample, we related plasma CD36 levels to a broad panel of adiposity measures, and observed a consistent and positive association of CD36 levels with VAT, SAT and BMI in different multivariable-adjusted statistical models. Furthermore, higher plasma CD36 levels have been related to greater liver fat content in clinical samples (e. g. in patients with NAFLD or chronic hepatitis C) (15,20,21), in individuals with obesity (17,18), and also in general population samples (22). In contrast to these former studies, plasma CD36 levels were unrelated to liver fat content in our analyses in a community-dwelling sample, free of apparent liver disease. One potential explanation for the lack of association in our study is that the association of CD36 with liver fat might be much more pronounced (or only apparent) in clinical settings or in obese individuals, but not in relatively healthy individuals from the community. Indeed, clinical study groups with significantly elevated plasma CD36 as previously reported had liver fat content of 20-30% (15,20,21), while our sample included many normal weight participants free of liver disease.

Furthermore, in our sample, CD36 levels were higher in individuals with NAFLD and elevated ALT concentrations as compared to individuals with NAFLD and normal ALT concentrations. Thus, CD36 concentrations might be more closely related to more severe liver cell damage (indicated by increased ALT levels) than to increased fat deposition in the liver.

In addition, differences in the definition of NAFLD and in the assessment of liver fat content have to be considered when comparing the results across studies (15,20-22). Previous studies among clinical samples used MRS (intrahepatic lipid content  $>5\%$ ) or liver biopsy in the absence of excessive alcohol intake to define NAFLD (15,20,21), while the current study defined NAFLD using MRI-measured LSI (log LSI  $\geq 3.0$ ) in the absence of high alcohol intake ( $\leq 7$  drinks/week for women and  $\leq 14$  drinks/week for men). In terms of liver fat content, a prior study conducted in a general population sample did not directly measure or visualize liver fat content, but estimated liver fat based on algorithms including BMI, WC, triglycerides, liver transaminases, fasting insulin and presence of diabetes and metabolic syndrome (22). Of note, most variables included in the algorithm to predict liver fat in the



previous study (22), such as BMI, WC, triglycerides and ALT, were also significantly correlated with plasma CD36 in our sample (**Supplemental Table S1**). By contrast, our study took advantage of MRI techniques to quantify liver fat content. However, liver fat content might not necessarily reflect NAFLD severity. A previous study suggested that liver fat content decreases in some patients with NAFLD progressing to cirrhosis (40). Long term prospective studies assessing fatty liver disease severity including fibrosis are warranted to better understand the association between circulating CD36 concentrations, liver fat (as assessed by MRI), biochemical markers of liver function and damage and NAFLD.

#### **Potential mechanisms for the observed associations**

Several experimental data support the observed association between plasma CD36 and adipose tissue traits. CD36 acts as a membrane fatty acid transporter (9) and has been shown to bind long-chain fatty acids (LCFA) and facilitate LCFA uptake across the plasma membrane in adipose tissue (41). In line with this biological function, an increased expression of CD36 has been observed on both, mRNA and plasma membrane level in VAT and SAT of obese individuals (42), and genetic variants in the *CD36* gene have been linked to lipid metabolism and obesity (10-12). Further, CD36 expression in the liver and plasma CD36 are correlated (15,20,21). Even though the mechanism of CD36 release into the circulation is currently unknown, it has been hypothesized that metabolic abnormalities such as inflammation in abdominal obesity could promote plasma CD36 release into the circulation (6).

#### **Strengths and limitations**

The strengths of the current study include its large sample size, the community-based design and the comprehensive clinical examination of the study participants, including assessment of different adipose tissues (including assessments of liver fat content on a continuous scale; a method that has been validated against MRS (24)) using MRI and detailed assessment of potentially confounding factors. Thus, we were able to adjust for a variety of lifestyle factors in our statistical analyses. The following limitations merit consideration. First, since this was a cross-sectional analysis, the causality and temporality of the relation of plasma CD36 levels with the different outcome measures could not be determined. Moreover, NAFLD was not diagnosed by liver biopsy in the current study, since this is not feasible and not ethically justifiable in a community-based setting. Therefore, we could not differentiate between different stages of NAFLD and their respective relations with plasma CD36 levels. Further studies are warranted that compare the accuracy of MRI-determined liver fat content to proton density fat-fraction, which is less confounded by biological, physical and technical factors (43) and has shown to accurately classify different grades of hepatic steatosis in patients with NAFLD (44,45). Additionally, the current study was based on a community-based sample from Northern Germany. Thus, our observations may not be generalizable to other populations.

#### **CONCLUSIONS**

In conclusion, higher plasma CD36 concentrations were statistically significantly associated with greater measures of general and abdominal adiposity, including higher values of MRI-determined SAT and VAT. However, CD36 levels were not related to MRI-determined liver fat content in our sample from the general population. These observations are consistent with the concept that plasma CD36 levels are relevant correlates of adipose tissue and provide initial evidence that plasma CD36 might be more closely related to SAT and VAT than liver fat content in the general population, a premise that needs further investigations. Individuals with NAFLD and elevated ALT had higher plasma CD36 levels than individuals with NAFLD and normal ATL levels. Thus, plasma CD36 may reflect more

advanced stages of NAFLD. Further population-based studies, e.g. with repeated measurements of plasma CD36 are warranted to describe trajectories of CD36 levels over time and their prospective association with metabolic and adiposity traits. Finally, studies on environmental, lifestyle and genetic factors associated with circulating CD36 are warranted.

### Acknowledgements:

We thank all the participants of the PopGen control cohort study. We also thank the technician Lone Larsen for dedicated measurements of CD36.

**Funding:** The current study is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC 22167-390884018 and by the German Federal Ministry of Education and Research (grant 01GR0468). The PopGen 2.0 network is supported by the German Federal Ministry of Education and Research (grant 01EY1103) and the Medical Faculty of the University of Kiel.

Deutsche Forschungsgemeinschaft <http://dx.doi.org/10.13039/501100001659>,

Germany's Excellence Strategy - EXC 22167-390884018, Not Applicable;

Bundesministerium für Bildung und Forschung

<http://dx.doi.org/10.13039/501100002347>, 01GR0468, Not Applicable;

Bundesministerium für Bildung und Forschung

<http://dx.doi.org/10.13039/501100002347>, 01EY1103, Not Applicable

Correspondence to: Manja Koch, Department of Nutrition, Harvard T.H. Chan School of Public Health, 655 Huntington Avenue, Boston, MA 02115. Phone: 617-432-1841, Fax: 617-432-2435, E-mail: [mkoch@hsph.harvard.edu](mailto:mkoch@hsph.harvard.edu).

Disclosure:

None.

### REFERENCES

1. World Health Organization. Obesity and overweight. 2018; <http://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>. Accessed 23 Feb 2018.
2. Fabbrini E, Magkos F. Hepatic Steatosis as a Marker of Metabolic Dysfunction. *Nutrients* 2015; 7:4995-5019
3. Abd El-Kader SM, El-Den Ashmawy EMS. Non-alcoholic fatty liver disease: The diagnosis and management. *World J Hepatol* 2015; 7:846-858
4. Pi-Sunyer X. The Medical Risks of Obesity. *Postgrad Med* 2009; 121:21-33
5. Vernon G, Baranova A, Younossi ZM. Systematic review: the epidemiology and natural history of non-alcoholic fatty liver disease and non-alcoholic steatohepatitis in adults. *Aliment Pharmacol Ther* 2011; 34:274-285
6. Koonen DP, Jensen MK, Handberg A. Soluble CD36- a marker of the (pathophysiological) role of CD36 in the metabolic syndrome? *Arch Physiol Biochem* 2011; 117:57-63
7. Miquilena-Colina ME, Lima-Cabello E, Sanchez-Campos S, Garcia-Mediavilla MV, Fernandez-Bermejo M, Lozano-Rodriguez T, Vargas-Castrillon J, Buque X, Ochoa B, Aspichueta P, Gonzalez-Gallego J, Garcia-Monzon C. Hepatic fatty acid translocase CD36 upregulation is associated with insulin resistance, hyperinsulinaemia and increased steatosis in non-alcoholic steatohepatitis and chronic hepatitis C. *Gut* 2011; 60:1394-1402
8. Greco D, Kotronen A, Westerbacka J, Puig O, Arkkila P, Kiviluoto T, Laitinen S, Kolak M, Fisher RM, Hamsten A, Auvinen P, Yki-Jarvinen H. Gene expression in human NAFLD. *Am J Physiol Gastrointest Liver Physiol* 2008; 294:G1281-1287

9. Bonen A, Tandon NN, Glatz JF, Luiken JJ, Heigenhauser GJ. The fatty acid transporter FAT/CD36 is upregulated in subcutaneous and visceral adipose tissues in human obesity and type 2 diabetes. *Int J Obes* 2006; 30:877-883
10. Lepretre F, Linton KJ, Lacquemant C, Vatin V, Samson C, Dina C, Chikri M, Ali S, Scherer P, Seron K, Vasseur F, Aitman T, Froguel P. Genetic study of the CD36 gene in a French diabetic population. *Diabetes Metab* 2004; 30:459-463
11. Bokor S, Legry V, Meirhaeghe A, Ruiz JR, Mauro B, Widhalm K, Manios Y, Amouyel P, Moreno LA, Molnar D, Dallongeville J, group HS. Single-nucleotide polymorphism of CD36 locus and obesity in European adolescents. *Obesity* 2010; 18:1398-1403
12. Heni M, Mussig K, Machicao F, Machann J, Schick F, Claussen CD, Stefan N, Fritsche A, Haring HU, Staiger H. Variants in the CD36 gene locus determine whole-body adiposity, but have no independent effect on insulin sensitivity. *Obesity* 2011; 19:1004-1009
13. Handberg A, Levin K, Hojlund K, Beck-Nielsen H. Identification of the oxidized low-density lipoprotein scavenger receptor CD36 in plasma: a novel marker of insulin resistance. *Circulation* 2006; 114:1169-1176
14. Handberg A, Skjelland M, Michelsen AE, Sagen EL, Krohg-Sorensen K, Russell D, Dahl A, Ueland T, Oie E, Aukrust P, Halvorsen B. Soluble CD36 in plasma is increased in patients with symptomatic atherosclerotic carotid plaques and is related to plaque instability. *Stroke* 2008; 39:3092-3095
15. Heeboll S, Poulsen MK, Ornstrup MJ, Kjaer TN, Pedersen SB, Nielsen S, Gronbaek H, Handberg A. Circulating sCD36 levels in patients with non-alcoholic fatty liver disease and controls. *Int J Obes* 2017; 41:262-267
16. Handberg A, Norberg M, Stenlund H, Hallmans G, Attermann J, Eriksson JW. Soluble CD36 (sCD36) clusters with markers of insulin resistance, and high sCD36 is associated with increased type 2 diabetes risk. *J Clin Endocrinol Metab* 2010; 95:1939-1946
17. Knosgaard L, Thomsen SB, Stockel M, Vestergaard H, Handberg A. Circulating sCD36 is associated with unhealthy fat distribution and elevated circulating triglycerides in morbidly obese individuals. *Nutr Diabetes* 2014; 4:e114
18. Knosgaard L, Kazankov K, Birkebaek NH, Holland-Fischer P, Lange A, Solvig J, Horlyck A, Kristensen K, Rittig S, Vilstrup H, Gronbaek H, Handberg A. Reduced sCD36 following weight loss corresponds to improved insulin sensitivity, dyslipidemia and liver fat in obese children. *Eur J Clin Nutr* 2016; 70:1073-1077
19. Glinborg D, Hojlund K, Andersen M, Henriksen JE, Beck-Nielsen H, Handberg A. Soluble CD36 and risk markers of insulin resistance and atherosclerosis are elevated in polycystic ovary syndrome and significantly reduced during pioglitazone treatment. *Diabetes Care* 2008; 31:328-334
20. Garcia-Monzon C, Lo Iacono O, Crespo J, Romero-Gomez M, Garcia-Samaniego J, Fernandez-Bermejo M, Dominguez-Diez A, Rodriguez de Cia J, Saez A, Porrero JL, Vargas-Castrillon J, Chavez-Jimenez E, Soto-Fernandez S, Diaz A, Gallego-Duran R, Madejon A, Miquilena-Colina ME. Increased soluble CD36 is linked to advanced steatosis in nonalcoholic fatty liver disease. *Eur J Clin Invest* 2014; 44:65-73
21. Petta S, Handberg A, Marchesini G, Camma C, Di Marco V, Cabibi D, Macaluso FS, Craxi A. High sCD36 plasma level is associated with steatosis and its severity in patients with genotype 1 chronic hepatitis C. *J Viral Hepat* 2013; 20:174-182
22. Handberg A, Hojlund K, Gastaldelli A, Flyvbjerg A, Dekker JM, Petrie J, Piatti P, Beck-Nielsen H. Plasma sCD36 is associated with markers of atherosclerosis, insulin resistance and fatty liver in a nondiabetic healthy population. *J Intern Med* 2012; 271:294-304

23. Mitra S, Fernandez-Del-Valle M, Hill JE. The role of MRI in understanding the underlying mechanisms in obesity associated diseases. *Biochim Biophys Acta Mol Basis Dis* 2017; 1863:1115-1131
24. Koch M, Borggrefe J, Barbaresko J, Groth G, Jacobs G, Siegert S, Lieb W, Muller MJ, Bosy-Westphal A, Heller M, Nothlings U. Dietary patterns associated with magnetic resonance imaging-determined liver fat content in a general population study. *Am J Clin Nutr* 2014; 99:369-377
25. The Human Protein Atlas. CD36. 2014; <https://www.proteinatlas.org/ENSG00000135218-CD36/tissue>. Accessed 10 March, 2018.
26. Nothlings U, Krawczak M. [PopGen. A population-based biobank with prospective follow-up of a control group]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2012; 55:831-835
27. Fischer K, Ruttgers D, Muller HP, Jacobs G, Kassubek J, Lieb W, Nothlings U. Association of Habitual Patterns and Types of Physical Activity and Inactivity with MRI-Determined Total Volumes of Visceral and Subcutaneous Abdominal Adipose Tissue in a General White Population. *PLoS One* 2015; 10:e0143925
28. di Giuseppe R, Koch M, Schlesinger S, Borggrefe J, Both M, Muller HP, Kassubek J, Jacobs G, Nothlings U, Lieb W. Circulating selenoprotein P levels in relation to MRI-derived body fat volumes, liver fat content, and metabolic disorders. *Obesity* 2017; 25:1128-1135
29. Nothlings U, Hoffmann K, Bergmann MM, Boeing H. Fitting portion sizes in a self-administered food frequency questionnaire. *J Nutr* 2007; 137:2781-2786
30. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR, Jr., Tudor-Locke C, Greer JL, Vezina J, Whitt-Glover MC, Leon AS. 2011 Compendium of Physical Activities: a second update of codes and MET values. *Med Sci Sports Exerc* 2011; 43:1575-1581
31. Barbaresko J, Siegert S, Koch M, Aits I, Lieb W, Nikolaus S, Laudes M, Jacobs G, Nothlings U. Comparison of two exploratory dietary patterns in association with the metabolic syndrome in a Northern German population. *Br J Nutr* 2014; 112:1364-1372
32. Fischer K, Moewes D, Koch M, Muller HP, Jacobs G, Kassubek J, Lieb W, Nothlings U. MRI-determined total volumes of visceral and subcutaneous abdominal and trunk adipose tissue are differentially and sex-dependently associated with patterns of estimated usual nutrient intake in a northern German population. *Am J Clin Nutr* 2015; 101:794-807
33. Taouli B, Ehman RL, Reeder SB. Advanced MRI methods for assessment of chronic liver disease. *AJR Am J Roentgenol* 2009; 193:14-27
34. Szczepaniak LS, Nurenberg P, Leonard D, Browning JD, Reingold JS, Grundy S, Hobbs HH, Dobbins RL. Magnetic resonance spectroscopy to measure hepatic triglyceride content: prevalence of hepatic steatosis in the general population. *Am J Physiol Endocrinol Metab* 2005; 288:E462-468
35. Long MT, Wang N, Larson MG, Mitchell GF, Palmisano J, Vasan RS, Hoffmann U, Speliotes EK, Vita JA, Benjamin EJ, Fox CS, Hamburg NM. Nonalcoholic fatty liver disease and vascular function: cross-sectional analysis in the Framingham heart study. *Arterioscler Thromb Vasc Biol* 2015; 35:1284-1291
36. National Institute on Alcohol Abuse and Alcoholism. Drinking Levels Defined. <https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/moderate-binge-drinking>. Accessed June 29, 2017.
37. Wang Y, Koch M, di Giuseppe R, Evans K, Borggrefe J, Nothlings U, Handberg A, K. Jensen M, Lieb W. Supplemental material for the paper "Association of plasma CD36 and body fat distribution". 2019. doi:10.6084/m9.figshare.8001962.v2.
38. Asakawa H, Tokunaga K, Kawakami F. Relationship of abdominal fat with metabolic disorders in diabetes mellitus patients. *Diabetes Res Clin Pract* 2002; 55:139-149



39. Kelley DE, Thaete FL, Troost F, Huwe T, Goodpaster BH. Subdivisions of subcutaneous abdominal adipose tissue and insulin resistance. *Am J Physiol Endocrinol Metab* 2000; 278:E941-948
40. Powell EE, Cooksley WG, Hanson R, Searle J, Halliday JW, Powell LW. The natural history of nonalcoholic steatohepatitis: a follow-up study of forty-two patients for up to 21 years. *Hepatology* 1990; 11:74-80
41. Glatz JF, Luiken JJ, Bonen A. Membrane fatty acid transporters as regulators of lipid metabolism: implications for metabolic disease. *Physiol Rev* 2010; 90:367-417
42. Grzegorzczak EA, Harasim-Symbor E, Lukaszuk B, Harasiuk D, Choromanska B, Mysliwiec P, Zendzian-Piotrowska M, Chabowski A. Lack of pronounced changes in the expression of fatty acid handling proteins in adipose tissue and plasma of morbidly obese humans. *Nutr Diabetes* 2018; 8:3
43. Reeder SB, Cruite I, Hamilton G, Sirlin CB. Quantitative Assessment of Liver Fat with Magnetic Resonance Imaging and Spectroscopy. *J Magn Reson Imaging* 2011; 34:spcone
44. Middleton MS, Heba ER, Hooker CA, Bashir MR, Fowler KJ, Sandrasegaran K, Brunt EM, Kleiner DE, Doo E, Van Natta ML, Lavine JE, Neuschwander-Tetri BA, Sanyal A, Loomba R, Sirlin CB. Agreement Between Magnetic Resonance Imaging Proton Density Fat Fraction Measurements and Pathologist-Assigned Steatosis Grades of Liver Biopsies From Adults With Nonalcoholic Steatohepatitis. *Gastroenterology* 2017; 153:753-761
45. Jayakumar S, Middleton MS, Lawitz EJ, Mantry PS, Caldwell SH, Arnold H, Mae Diehl A, Ghalib R, Elkhatab M, Abdelmalek MF, Kowdley KV, Stephen Djedjos C, Xu R, Han L, Mani Subramanian G, Myers RP, Goodman ZD, Afdhal NH, Charlton MR, Sirlin CB, Loomba R. Longitudinal correlations between MRE, MRI-PDFF, and liver histology in patients with non-alcoholic steatohepatitis: Analysis of data from a phase II trial of selonsertib. *J Hepatol* 2019; 70:133-141

**Table 1.** Characteristics of the study sample (n=575), stratified by sex-specific quartiles of plasma CD36.

	Quartiles of CD36			
	Q1 (n=143)	Q2 (n=143)	Q3 (n=145)	Q4 (n=144)
Median CD36, arbitrary unit (Q25, Q75) for men	0.57 (0.48, 0.70)	1.02 (0.89, 1.13)	1.53 (1.38, 1.64)	2.47 (1.99, 3.02)
Median CD36, arbitrary unit (Q25, Q75) for women	0.54 (0.43, 0.63)	0.89 (0.81, 1.02)	1.23 (1.14, 1.40)	1.95 (1.81, 2.34)
Men, (%)	57.3	58.0	56.6	57.6
Age, y	62.4 (53.7, 70.9)	64.0 (54.7, 72.4)	61.7 (55.5, 69.4)	61.3 (53.5, 69.0)
Body mass index, (kg/m <sup>2</sup> )	26.4 ± 4.0	26.8 ± 4.4	27.1 ± 4.1	27.9 ± 4.6
Waist circumference for men, cm	97.3 ± 9.9	99.6 ± 10.3	99.9 ± 10.0	101.2 ± 11.2
Waist circumference for women, cm	88.1 ± 12.6	90.0 ± 12.6	89.9 ± 13.3	93.5 ± 14.1
Systolic blood pressure, mmHg	138.3 ± 17.2	139.5 ± 18.5	140.8 ± 19.2	138.9 ± 18.8
Prevalent diabetes, %	8.4	13.3	8.28	13.2
Current smokers, %	11.9	11.9	6.90	9.72
Physical activity, MET-hr/wk	97.5 (56.8, 141.7)	92.8 (58.0, 123.5)	84.5 (61.8, 124.8)	88.0 (56.3, 128.8)
High education, %	39.2	34.3	34.5	32.6
Alcohol consumption, g/d	9.03 (2.89, 15.9)	10.1 (2.62, 21.2)	9.15 (3.68, 19.5)	9.93 (4.91, 17.0)



SAT, dm <sup>3</sup>	5.40 (4.36, 7.53)	6.20 (4.46, 8.11)	6.12 (4.74, 8.18)	6.84 (5.28, 9.38)
VAT, dm <sup>3</sup>	3.09 (2.12, 4.64)	4.10 (2.56, 5.44)	3.78 (2.41, 4.96)	4.21 (2.89, 5.62)
ALT, IU/L	21.0 (16.0, 27.0)	22.0 (17.0, 28.0)	23.0 (17.0, 31.0)	25.0 (18.0, 32.0)
Liver signal intensity	2.89 ± 0.41	2.93 ± 0.39	2.98 ± 0.43	2.95 ± 0.54
Fatty liver disease, %	37.8	37.8	42.8	43.8
Non-alcoholic fatty liver disease, %	32.6	32.1	37.1	37.7

Data are reported as mean ± standard deviation for continuous normally distributed variables (body mass index, waist circumference, systolic blood pressure and liver signal intensity) and as median (Q25, Q75) for continuous skewed distributed variables (CD36, age, physical activity, alcohol consumption, SAT, VAT and ALT) and as percentage for categorical variables. **Abbreviations:** CD36, cluster of differentiation 36; MET, metabolic equivalent; SAT, subcutaneous abdominal adipose tissue; VAT, visceral abdominal adipose tissue; ALT, alanine aminotransferase.

**Table 2.** Multivariable-adjusted means (95% CI) of different MRI-derived adiposity traits and measures of liver fat, stratified by sex-specific quartiles of plasma CD36 (*n*=575).

	Multivariable-adjusted means (95% CI) by quartiles of CD36				<i>P</i> for trend <sup>a</sup>
	Q1 ( <i>n</i> =143)	Q2 ( <i>n</i> =147)	Q3 ( <i>n</i> =140)	Q4 ( <i>n</i> =146)	
<i>Liver signal intensity</i>					
Model 1 <sup>b</sup>	2.90 (2.83, 2.98)	2.93 (2.86, 3.00)	2.99 (2.91, 3.06)	2.96 (2.89, 3.03)	0.38
Model 2 <sup>c</sup>	3.05 (2.96, 3.14)	3.04 (2.95, 3.13)	3.11 (3.02, 3.20)	3.07 (2.98, 3.15)	0.36
Mean difference (95% CI), Model 2	0 (Reference)	-0.01 (-0.12, 0.11)	0.06 (-0.05, 0.18)	0.02 (-0.10, 0.14)	
<i>BMI, kg/m<sup>2</sup></i>					
Model 1 <sup>b</sup>	26.4 (25.7, 27.1)	26.8 (26.1, 27.5)	27.1 (26.4, 27.8)	27.9 (27.2, 28.6)	0.022
Model 2 <sup>c</sup>	27.8 (26.9, 28.6)	27.9 (27.1, 28.8)	28.2 (27.3, 29.1)	28.9 (28.0, 29.7)	0.013
Mean difference (95% CI), Model 2	0 (Reference)	0.16 (-0.99, 1.31)	0.47 (-0.67, 1.62)	1.13 (-0.03, 2.28)	
<i>VAT, dm<sup>3</sup></i>					
Model 1 <sup>b</sup>	3.38 (3.08, 3.68)	4.07 (3.77, 4.36)	3.86 (3.56, 4.16)	4.22 (3.93, 4.52)	<0.001
Model 2 <sup>c</sup>	4.13 (3.77, 4.48)	4.65 (4.31, 5.00)	4.50 (4.14, 4.86)	4.71 (4.36, 5.06)	<0.001
Mean difference (95% CI), Model 2	0 (Reference)	0.53 (0.05, 1.00)	0.38 (-0.10, 0.86)	0.59 (0.11, 1.07)	
<i>SAT, dm<sup>3</sup></i>					
Model 1 <sup>b</sup>	6.40 (5.85, 6.94)	6.99 (6.46, 7.53)	7.10 (6.55, 7.65)	7.87 (7.33, 8.41)	0.002
Model 2 <sup>c</sup>	7.61 (6.96, 8.27)	7.99 (7.36, 8.63)	8.21 (7.54, 8.87)	8.74 (8.10, 9.38)	<0.001
Mean difference (95% CI), Model 2	0 (Reference)	0.38 (-0.49, 1.25)	0.59 (-0.29, 1.47)	1.13 (0.25, 2.01)	

<sup>a</sup>Linear trend was tested per standard deviation increment of plasma CD36 levels. <sup>b</sup>Model 1: Adjusted for age and sex. <sup>c</sup>Model 2: Model 1 additionally adjusted for education levels ( $\leq 9$ , 10 or  $\geq 11$  years), smoking status (never, former or current smoker), physical activity (continuous), prevalent diabetes (yes or no) and alcohol intake (continuous). **Abbreviations:** BMI, body mass index; CD36, cluster of differentiation 36; CI, confidence interval; VAT, visceral abdominal adipose tissue; SAT, subcutaneous abdominal adipose tissue; Q, quartile.